

JPG 32, 1025 (2006)

NPBPS 186, 207 (2009)

HADRONIC EFFECTS IN LOW-ENERGY QCD:
ADLER FUNCTION AND τ DECAY

A.V. Nesterenko


Bogoliubov Laboratory of Theoretical Physics

Joint Institute for Nuclear Research, Dubna, 141980, Russia

Fourteenth Lomonosov Conference on Elementary Particle Physics
Moscow, Russia, 19 – 25 August 2009

INTRODUCTION

Hadronic vacuum polarization function $\Pi(q^2)$ plays a crucial role in various issues of elementary particle physics. Indeed, the theoretical description of some strong interaction processes and of the hadronic contributions to electroweak observables is inherently based on $\Pi(q^2)$:



- electron–positron annihilation into hadrons
- hadronic τ lepton decay
- muon anomalous magnetic moment
- running of the electromagnetic coupling

The cross-section of e^+e^- annihilation into hadrons reads

$$\sigma = 4\pi^2 \frac{2\alpha^2}{s^3} L^{\mu\nu} \Delta_{\mu\nu},$$


where $s = q^2 = (p_1 + p_2)^2 > 0$,

$$L_{\mu\nu} = \frac{1}{2} \left[q_\mu q_\nu - g_{\mu\nu} q^2 - (p_1 - p_2)_\mu (p_1 - p_2)_\nu \right],$$

$$\Delta_{\mu\nu} = (2\pi)^4 \sum_{\Gamma} \delta(p_1 + p_2 - p_\Gamma) \langle 0 | J_\mu(-q) | \Gamma \rangle \langle \Gamma | J_\nu(q) | 0 \rangle,$$

Γ denotes a final hadron state, and $J_\mu = \sum_f Q_f : \bar{q} \gamma_\mu q :$ stands for the electromagnetic quark current.

It is worth stressing that $\Delta_{\mu\nu}(q^2)$ exists only for $q^2 \geq 4m_\pi^2$, since otherwise no hadron state Γ could be excited:



■ *R.P.Feynman (1972); S.L.Adler, PRD10 (1974).*

The hadronic tensor can be represented as $\Delta_{\mu\nu} = 2 \text{Im} \Pi_{\mu\nu}$,
 $\Pi_{\mu\nu}(q^2) = i \int e^{iqx} \langle 0 | T \{ J_\mu(x) J_\nu(0) \} | 0 \rangle d^4x = (q_\mu q_\nu - g_{\mu\nu} q^2) \Pi(q^2)$.

The hadronic vacuum polarization function $\Pi(q^2)$ satisfies the once-subtracted dispersion relation (cut for $q^2 \geq 4m_\pi^2$)

$$\Pi(q^2) = \Pi(q_0^2) - (q^2 - q_0^2) \int_{4m_\pi^2}^{\infty} \frac{R(s)}{(s - q^2)(s - q_0^2)} ds,$$

where $m_\pi \simeq 135 \text{ MeV}$ is the mass of the π meson and $R(s)$ denotes the measurable ratio of two cross-sections:

$$R(s) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0_+} \left[\Pi(s - i\varepsilon) - \Pi(s + i\varepsilon) \right] = \frac{\sigma(e^+e^- \rightarrow \text{hadrons}; s)}{\sigma(e^+e^- \rightarrow \mu^+\mu^-; s)}.$$

It is worth noting here that $R(s) \equiv 0$ for $s < 4m_\pi^2$ because of the kinematic restrictions mentioned above:

■ *R.P.Feynman (1972).*

For practical purposes it proves to be convenient to deal with the so-called Adler function $D(Q^2)$ ($Q^2 = -q^2 \geq 0$):

$$D(Q^2) = \frac{d \Pi(-Q^2)}{d \ln Q^2}, \quad D(Q^2) = Q^2 \int_{4m_\pi^2}^{\infty} \frac{R(s)}{(s + Q^2)^2} ds,$$


which plays an indispensable role for the congruous analysis of the timelike and spacelike experimental data:

■ *S.L.Adler PRD10 (1974); A.Rujula, H.Georgi PRD13 (1976); J.D.Bjorken (1989).*

The inverse relation between

$D(Q^2)$ and $R(s)$ reads


$$R(s) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_{s+i\varepsilon}^{s-i\varepsilon} D(-\zeta) \frac{d\zeta}{\zeta}$$



■ *A.V.Radyushkin (1982), hep-ph/9907228;*
N.V.Krasnikov, A.A.Pivovarov PLB116 (1982).

Although there are no direct measurements of the Adler function, it can be restored by employing the experimental data on $R(s)$ (overall factor $N_c \sum_f Q_f^2$ is omitted):

$$D_{\text{exp}}(Q^2) = Q^2 \int_{4m_\pi^2}^{s_0} \frac{R_{\text{exp}}(s)}{(s + Q^2)^2} ds + Q^2 \int_{s_0}^{\infty} \frac{R_{\text{theor}}(s)}{(s + Q^2)^2} ds$$




There is also a number of lattice simulations, which generally agree with the shown result:

■ *JLQCD and TWQCD Collabs., PRD79 (2009); QCDSF Collab., NPB688 (2004).*

On the one hand, perturbation theory provides an explicit expression for the Adler function valid at $Q^2 \rightarrow \infty$:

$$D_{\text{pert}}^{(\ell)}(Q^2) = 1 + \sum_{j=1}^{\ell} d_j \left[\alpha_s^{(\ell)}(Q^2) \right]^j.$$



■ *S.G.Gorishny, A.L.Kataev, S.A.Larin PLB259 (1991); L.R.Surguladze, M.A.Samuel PRL66 (1991); P.A.Baikov, K.G.Chetyrkin, J.H.Kuhn, PRL101 (2008).*

On the other hand, this perturbative approximation is inconsistent with the dispersion relation for $D(Q^2)$ due to unphysical singularities of the strong running coupling $\alpha_s(Q^2)$:

$$D_{\text{pert}}^{(1)}(Q^2) = 1 + d_1 \alpha_s^{(1)}(Q^2), \quad \alpha_s^{(1)}(Q^2) = \frac{4\pi}{\beta_0} \frac{1}{\ln(Q^2/\Lambda^2)},$$

where $d_1 = 1/\pi$ and $\beta_0 = 11 - 2n_f/3$.

Dispersion relation imposes stringent constraints on $D(Q^2)$:
$$D(Q^2) = Q^2 \int_{4m_\pi^2}^{\infty} \frac{R(s)}{(s + Q^2)^2} ds$$

- ⊙ Since $R(s)$ assumes finite values and $R(s) \rightarrow \text{const}$ when $s \rightarrow \infty$, then $D(Q^2) \rightarrow 0$ at $Q^2 \rightarrow 0$ (holds for $m_\pi \neq 0$ only)
- ⊙ Adler function possesses the only cut $Q^2 \leq -4m_\pi^2$ along the negative semiaxis of real Q^2


PRIMARY OBJECTIVE: to merge these nonperturbative constraints with perturbative result for the Adler function.

NEW INTEGRAL REPRESENTATION FOR $D(Q^2)$

This objective can be achieved by deriving the integral representations for the Adler function and $R(s)$ -ratio, which involve a common spectral function.

$$D(Q^2) = Q^2 \int_{4m_\pi^2}^{\infty} \frac{R(s)}{(s + Q^2)^2} ds$$

$$R(s) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_{s+i\varepsilon}^{s-i\varepsilon} D(-\zeta) \frac{d\zeta}{\zeta}$$



Parton model prediction + kinematic restriction on $R(s)$:

$$R_0(s) = \theta(s - 4m_\pi^2)$$



$$D_0(Q^2) = \frac{Q^2}{Q^2 + 4m_\pi^2}$$

■ *R.P. Feynman (1972).*

$$D(Q^2) = \frac{Q^2}{Q^2 + 4m_\pi^2} + d(Q^2)$$

$$R(s) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0_+} \int_{s+i\varepsilon}^{s-i\varepsilon} D(-\zeta) \frac{d\zeta}{\zeta}$$

$$R(s) = \theta(s - 4m_\pi^2) \left[1 + \int_s^\infty \rho_D(\sigma) \frac{d\sigma}{\sigma} \right]$$

$$D(Q^2) = Q^2 \int_{4m_\pi^2}^\infty \frac{R(s)}{(s + Q^2)^2} ds$$

$$D(Q^2) = \frac{Q^2}{Q^2 + 4m_\pi^2} \left[1 + \int_{4m_\pi^2}^\infty \rho_D(\sigma) \frac{\sigma - 4m_\pi^2}{\sigma + Q^2} \frac{d\sigma}{\sigma} \right]$$

$$\rho_D(\sigma) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0_+} \left[D_{\text{theor}}(-\sigma - i\varepsilon) - D_{\text{theor}}(-\sigma + i\varepsilon) \right] = - \frac{d R_{\text{exp}}(\sigma)}{d \ln \sigma}$$

■ A.V. Nesterenko, J. Papavassiliou, JPG32 (2006).

Thus one arrives at the following integral representations:

$$D(Q^2) = \frac{Q^2}{Q^2 + 4m_\pi^2} \left[1 + \int_{4m_\pi^2}^{\infty} \rho_D(\sigma) \frac{\sigma - 4m_\pi^2}{\sigma + Q^2} \frac{d\sigma}{\sigma} \right]$$

$$R(s) = \theta(s - 4m_\pi^2) \left[1 + \int_s^{\infty} \rho_D(\sigma) \frac{d\sigma}{\sigma} \right]$$

$$\rho_D(\sigma) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0_+} \left[D_{\text{theor}}(-\sigma - i\varepsilon) - D_{\text{theor}}(-\sigma + i\varepsilon) \right] = - \frac{d R_{\text{exp}}(\sigma)}{d \ln \sigma}$$

■ *A.V. Nesterenko, J. Papavassiliou, JPG32 (2006).*

- all nonperturbative constraints on $D(Q^2)$ are satisfied
- congruent analysis of spacelike and timelike processes

In the limit $m_\pi = 0$ the obtained expressions become identical to those of the Analytic perturbation theory

■ *D.V. Shirkov, I.L. Solovtsov, PRL79 (1997); EPJC22 (2001); TMP150 (2007).*


There is no unique way to compute the corresponding spectral density by making use of perturbative $D_{\text{pert}}(Q^2)$. In what follows the one-loop spectral function is adopted:

$$\rho^{(1)}(\sigma) = \left(1 + \frac{1}{\sigma}\right) \frac{1}{\ln^2 \sigma + \pi^2}$$

■ *A.V. Nesterenko, PRD62 (2000); PRD64 (2001).*

ADVANTAGES :

- unphysical perturbative singularities are eliminated
- additional parameters are not introduced
- reasonable agreement with $D_{\text{exp}}(Q^2)$ for all energies



INCLUSIVE τ LEPTON DECAY

The inclusive semileptonic branching ratio:

$$R_\tau = \frac{\Gamma(\tau^- \rightarrow \text{hadrons}^- \nu_\tau)}{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = R_{\tau,V} + R_{\tau,A} + R_{\tau,S}.$$

Its nonstrange part associated with vector quark currents:

$$R_{\tau,V} = \frac{N_c}{2} |V_{ud}|^2 S_{EW} (\Delta_{\text{QCD}} + \delta'_{EW}) = 1.764 \pm 0.016$$

■ *OPAL Collab., EPJC7 (1999); ALEPH Collab., EPJC4 (1998), RMP78 (2006).*

In this equation $N_c = 3$, $|V_{ud}| = 0.9738 \pm 0.0005$, $\delta'_{EW} = 0.0010$,
 $S_{EW} = 1.0194 \pm 0.0050$, $M_\tau = 1.777 \text{ GeV}$, and

$$\Delta_{\text{QCD}} = 2 \int_0^{M_\tau^2} \left(1 - \frac{s}{M_\tau^2}\right)^2 \left(1 + 2 \frac{s}{M_\tau^2}\right) R(s) \frac{ds}{M_\tau^2}.$$

Perturbative approach:

$$\Delta_{\text{QCD}} = 1 + d_1 \alpha_s^{(1)}(M_\tau^2) \longrightarrow \Lambda = (678 \pm 55) \text{ MeV}, \quad n_f = 2$$

■ *E. Braaten, S. Narison, A. Pich, NPB373 (1992).*

Current analysis:

$$\Delta_{\text{QCD}} = 1 + d_1 \alpha_{\text{TL}}^{(1)}(M_\tau^2) - \delta_\Gamma + \frac{4}{\beta_0} \int_\chi^1 f(\xi) \rho^{(1)}(\xi M_\tau^2) d\xi - d_1 \delta_\Gamma \alpha_{\text{TL}}^{(1)}(m_\Gamma^2),$$

$$f(\xi) = \xi^3 - 2\xi^2 + 2, \quad \chi = \frac{m_\Gamma^2}{M_\tau^2}, \quad \delta_\Gamma = \chi f(\chi) \simeq 0.048, \quad d_1 = \frac{1}{\pi},$$

$$\alpha_{\text{TL}}^{(1)}(s) = \frac{4\pi}{\beta_0} \theta(s - m_\Gamma^2) \int_s^\infty \rho^{(1)}(\sigma) \frac{d\sigma}{\sigma}, \quad m_\Gamma = m_{\pi^0} + m_{\pi^-}$$

- **massive case:** $\Lambda = (941 \pm 86) \text{ MeV}$
- **massless limit:** $\Lambda = (493 \pm 56) \text{ MeV}$

■ *A.V. Nesterenko, NPBPS186 (2009).*

SUMMARY

- New integral representations for the Adler function and $R(s)$ -ratio are derived
- These representations possess appealing features:
 - unphysical perturbative singularities are eliminated
 - additional parameters are not introduced
 - the π^2 -terms are automatically taken into account
 - reasonable description of $D(Q^2)$ in entire energy range
- The effects due to the pion mass play a substantial role in the analysis of the inclusive τ lepton hadronic decay