

THE DIPOLE POLARIZABILITY OF THE DOUBLY MAGIC NUCLEI

Arsenyev Nikolay

Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna, Russia

In collaboration with

A. P. Severyukhin — BLTP, JINR, Dubna

Seminar

in memory of Prof. B. S. Ishkanov

“Photonuclear research: status and prospects”

Skobeltsyn Institute of Nuclear Physics MSU

Moscow, Russia

26 October 2023

Outline

Introduction

Realization of QRPA with the Skyrme EDF

Results and discussion


Conclusions

N. Arsenyev



Introduction

$E1$ strength in (spherical) atomic nuclei




Courtesy: N. Pietralla

N. Arsenyev



Introduction

The electric dipole polarizability



External Field



Balanced



Displacement
Dipole Polarizability

Restoring Force

Courtesy: A. Tamii

$$\alpha_D = \frac{8\pi}{9} \sum_{\nu} \frac{B(E1; 0_{g.st.}^+ \rightarrow 1_{\nu}^-)}{E_{1_{\nu}^-}}$$

N. Arsenyev



REALIZATION OF QRPA

N. Arsenyev



Realization of QRPA

The starting point of the method is **the HF-BCS calculations** of the ground state, where spherical symmetry is assumed for the ground states. The continuous part of the single-particle spectrum is discretized by diagonalizing the HF Hamiltonian on a harmonic oscillator basis.

J. P. Blaizot and D. Gogny, Nucl. Phys. A284, 429 (1977).

The residual interaction in the particle-hole channel V_{res}^{ph} and in the particle-particle channel V_{res}^{pp} can be obtained as the second derivative of the energy density functional \mathcal{H} with respect to the particle density ρ and the pair density $\tilde{\rho}$, respectively.

$$V_{res}^{ph} \sim \frac{\delta^2 \mathcal{H}}{\delta \rho_1 \delta \rho_2} \quad V_{res}^{pp} \sim \frac{\delta^2 \mathcal{H}}{\delta \tilde{\rho}_1 \delta \tilde{\rho}_2}.$$

G. T. Bertsch and S. F. Tsai, Phys. Rep. 18, 125 (1975).

N. Arsenyev



Realization of QRPA

We employ the effective Skyrme interaction in the particle-hole channel

$$V(\vec{r}_1, \vec{r}_2) = t_0 \left(1 + x_0 \hat{P}_\sigma\right) \delta(\vec{r}_1 - \vec{r}_2) + \frac{t_1}{2} \left(1 + x_1 \hat{P}_\sigma\right) \left[\delta(\vec{r}_1 - \vec{r}_2) \vec{k}^2 + \vec{k}'^2 \delta(\vec{r}_1 - \vec{r}_2) \right] \\ + t_2 \left(1 + x_2 \hat{P}_\sigma\right) \vec{k}' \cdot \delta(\vec{r}_1 - \vec{r}_2) \vec{k} + \frac{t_3}{6} \left(1 + x_3 \hat{P}_\sigma\right) \delta(\vec{r}_1 - \vec{r}_2) \rho^\alpha \left(\frac{\vec{r}_1 + \vec{r}_2}{2} \right) \\ + iW_0 (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot \left[\vec{k}' \times \delta(\vec{r}_1 - \vec{r}_2) \vec{k} \right].$$

T. H. R. Skyrme, Nucl. Phys. 9, 615 (1959).

D. Vautherin and D. M. Brink, Phys. Rev. C5, 626 (1972).

The Hamiltonian includes the pairing correlations are generated by the density-dependent zero-range force in the particle-particle channel

$$V_{pair}(\vec{r}_1, \vec{r}_2) = V_0 \left(1 - \eta \frac{\rho(r_1)}{\rho_{sat}} \right) \delta(\vec{r}_1 - \vec{r}_2),$$

where ρ_{sat} is the nuclear saturation density; η and V_0 are model parameters. For example, $\eta=0$ and $\eta=1$ are the case of a volume interaction and a surface-peaked interaction, respectively.

N. Arsenyev



Realization of QRPA

We introduce the phonon creation operators

$$Q_{\lambda\mu i}^+ = \frac{1}{2} \sum_{jj'} \left[X_{jj'}^{\lambda i} A^+(jj'; \lambda\mu) - (-1)^{\lambda-\mu} Y_{jj'}^{\lambda i} A(jj'; \lambda - \mu) \right],$$
$$A^+(jj'; \lambda\mu) = \sum_{mm'} C_{jmj'm'}^{\lambda\mu} \alpha_{jm}^+ \alpha_{j'm'}^+.$$

The index λ denotes total angular momentum and μ is its z-projection in the laboratory system. One assumes that the ground state is the QRPA phonon vacuum $|0\rangle$ and one-phonon excited states are $Q_{\lambda\mu i}^+|0\rangle$ with the normalization condition

$$\langle 0|[Q_{\lambda\mu i}, Q_{\lambda\mu i'}^+]|0\rangle = \delta_{ii'}.$$

Making use of the linearized equation-of-motion approach one can get the QRPA equations

$$\begin{pmatrix} \mathcal{A} & \mathcal{B} \\ -\mathcal{B} & -\mathcal{A} \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix} = \omega \begin{pmatrix} X \\ Y \end{pmatrix}.$$

Solutions of this set of linear equations yield the one-phonon energies ω and the amplitudes X, Y of the excited states.

RESULTS AND DISCUSSION

Details of calculations

The dipole transition probabilities can be expressed as

$$B(E1; 0_{gs}^+ \rightarrow 1_i^-) = \left| e_{eff}^{(n)} \langle i | \hat{M}^{(n)} | 0 \rangle + e_{eff}^{(p)} \langle i | \hat{M}^{(p)} | 0 \rangle \right|^2,$$

where $\hat{M}^{(p)} = \sum_i^Z r_i Y_{1\mu}(\hat{r}_i)$ and $\hat{M}^{(n)} = \sum_i^N r_i Y_{1\mu}(\hat{r}_i)$. The spurious 1^- state is excluded from the excitation spectra by introduction of the effective neutron $e_{eff}^{(n)} = -Z/A e$ and proton $e_{eff}^{(p)} = N/A e$ charges.

A. Bohr and B. Mottelson, Nuclear Structure Vol. II (Benjamin, New York 1975).


For our systematic analysis of the electric dipole polarizability, we employ 21 Skyrme interactions: LNS, SAMi, SGII, SIII, SK255, SkI2, SkI3, SkI5, SkM*, SkP, SkT5, SkT6, SkT7, SkX, SLy4, SLy5, SVbas, SVmas07, SVmas08, SVmas10, and SVmin. The choice of these parameterizations is due to the large range of values for the effective nucleon mass $m^* = 0.58-1.00$ and the symmetry energy at saturation density $J = 26.8-37.4$ MeV.

M. Dutra et al., Phys. Rev. C85, 035201 (2012).

N. Arsenyev



The photoabsorption cross section for ^{40}Ca




N. N. Arsenyev, A. P. Severyukhin, in preparation.

R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

N. Arsenyev



The photoabsorption cross section for ^{48}Ca




N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

J. Birkhan et al., PRL 118, 252501 (2017).

N. Arsenyev



The electric dipole polarizability α_D in $^{40,48}\text{Ca}$



N. N. Arsenyev, A. P. Severyukhin, in preparation.

J. Birkhan et al., PRL 118, 252501 (2017).


R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

G. Hagen et al., Nature Phys. 12, 186 (2016).

N. Arsenyev



The electric dipole polarizability α_D : Expt. vs Theory




N. N. Arsenyev, A. P. Severyukhin, in preparation.

N. Arsenyev



The neutron skin thickness ΔR_{np} in $^{40,48}\text{Ca}$



N. N. Arsenyev, A. P. Severyukhin, in preparation.

J. Birkhan et al., PRL 118, 252501 (2017).


R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

G. Hagen et al., Nature Phys. 12, 186 (2016).

N. Arsenyev



The neutron skin thickness ΔR_{np} in ^{132}Sn and ^{208}Pb



N. N. Arsenyev, A. P. Severyukhin, in preparation.

A. Tamii et al., PRL 107, 062502 (2011).

A. Klimkiewicz et al., Phys. Rev. C76, 051603 (2007).

K. Hebeler et al., PRL 105, 161102 (2010).

N. Arsenyev



Pearson correlation coefficient

In statistics, the Pearson correlation coefficient (r) is a measure of linear correlation between two sets of data: $(a_1, b_1), \dots, (a_N, b_N)$. The Pearson coefficient calculate as

$$r = \frac{\sum_{i=1}^N (a_i - \bar{a})(b_i - \bar{b})}{\sigma_a \sigma_b},$$

where \bar{a} , \bar{b} represent the mean values and σ_a , σ_b the root mean square of the sample. The coefficient r always has a value between -1 and $+1$. Correlations equal to -1 or $+1$ correspond to data points lying exactly on a line. In fact, we may transform a_k to \tilde{a}_k , where

$$\tilde{a}_k = r \frac{\sigma_a}{\sigma_b} (b_k - \bar{b}) + \bar{a}.$$





For any value of the coefficient r , the value of \tilde{a}_k is the best linear approximation for a_k .

N. Draper and H. Smith, Applied Regression Analysis (Wiley, New York, 1998).

N. Arsenyev



Correlations: α_D vs ΔR_{np}



N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

N. Arsenyev



The nuclear symmetry energy J

One of the key properties of nuclear matter is the symmetry energy (J), which is particularly important in modeling nuclear matter and finite nuclei because it probes the isospin part of the Skyrme interaction. It is defined as

$$J(\rho) = \frac{\hbar^2}{6m} \left(\frac{3\pi^2}{2} \right)^{2/3} \rho^{2/3} - \frac{1}{8} t_0 (2x_0 + 1) \rho - \frac{1}{24} \left(\frac{3\pi^2}{2} \right)^{2/3} \rho^{2/3} [3t_1 x_1 - t_2 (4 + 5x_2)] - \frac{1}{48} t_3 (2x_3 + 1) \rho^{\alpha+1}.$$

Saturation properties of a few Skyrme parametrizations used in this work.





	SLy4	SAMi	SIII	SGII	SkM*	SkP	SkI2	SkT5	SK255	LNS
J	32.00	28.00	28.16	26.83	30.03	30.00	33.37	37.00	37.40	33.43
ρ_{sat}	0.160	0.159	0.145	0.158	0.160	0.163	0.158	0.164	0.157	0.175

M. Dutra et al., Phys. Rev. C85, 035201 (2012).

N. Arsenyev



Correlations: α_D vs ΔR_{np} and J




N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

N. Arsenyev



Estimation of the symmetry energy J

We carried out a theoretical analysis of the recently measured α_D and ΔR_{np} in ^{48}Ca , ^{68}Ni , and ^{208}Pb to extract information about the symmetry energy, by using a strong correlation between $\alpha_D J$ and ΔR_{np} . Combining the experimental data and the RPA theory constraints yields the interval of $J = 30 - 37 \text{ MeV}^\dagger$.



N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

A. Tamii et al., PRL 107, 062502 (2011).

D. M. Rossi et al., PRL 111, 242503 (2013).

J. Birkhan et al., Phys. Rev. Lett. 118, 252501 (2017).

[†] The current global average $J = 31.60 \pm 0.92 \pm 2.66 \text{ MeV}$ obtained using the 28 analyses of terrestrial nuclear experiments and astrophysical observations [B. A. Li and X. Han, Phys. Lett. B727, 276 (2013)]

N. Arsenyev



Conclusions

The Skyrme RPA calculations have been performed for the electric dipole response in the magic nuclei ^{16}O , $^{40,48,60}\text{Ca}$, $^{56,68,78}\text{Ni}$, ^{90}Zr , $^{100,132}\text{Sn}$ and ^{208}Pb . We have computed the nuclear dipole polarizability (α_{D}) and neutron skin thickness (ΔR_{np}) of these nuclei using a broad set of Skyrme functionals. It is shown that the neutron skin thickness is correlated the product of the electric dipole polarizability and the symmetry energy (J) at saturation density.

We carried out a theoretical analysis of the recently measured α_{D} and ΔR_{np} in ^{48}Ca , ^{68}Ni , and ^{208}Pb to extract information about the symmetry energy, by using a strong correlation between α_{D} and ΔR_{np} . Combining the experimental data and the RPA theory (using a broad set of Skyrme functionals) constraints yields the interval of $J=30-37$ MeV.

This work was partly supported by the Russian Science Foundation (Grant No. RSF-21-12-00061) and the National Research Foundation of South Africa (Grant No. 129603).

N. Arsenyev



THE END

N. Arsenyev

