



Проблемы холодной темной материи как главного гравитирующего вещества


Сильченко О.К., ГАИШ МГУ

Эволюция крупномасштабной структуры: темная материя - **ХОЛОДНАЯ**



- Основной механизм — иерархическое сгущивание материи под действием гравитационной неустойчивости
- Подбор параметров — чтобы получить правильную конечную картину к моменту $z=0$

Карликовые галактики на малых z : Comparison of the distribution function of line widths V_{los} for galaxies in the Local Volume with theoretical predictions for the Λ CDM(left-hand panel) and the WDM models (right-hand panel, Planck cosmological parameters).



Anatoly Klypin et al. MNRAS 2015;454:1798-1810

Сравнение количества галактик разных масс на разных z с моделями




Figure 16. A comparison between our data and the models from the Millennium Simulation (e.g. De Lucia et al. 2006). The data shown are the same for the massive galaxy sample plotted in Fig. 4. The dashed line shows the predicted evolution in the number and stellar mass densities for $10^{11} M_\odot < M_* < 10^{11.5} M_\odot$ systems, while the solid line shows the same predictions for galaxies with stellar masses $M_* > 10^{11.5} M_\odot$. As can be seen, for the most part the simulated massive galaxies do not assemble quickly enough to match the observations.

Downsizing!

Perez-Gonzalez et al. 2008

Serjeant et al. 2008




FIG. 6.— Fraction of the local stellar mass density at a given redshift for several mass intervals (referring to more massive systems). Only results for our 75% completeness level at each redshift are shown.




Figure 12. Left: SFR per unit galaxy mass, $(1/M_*) dM_*/dt$. This has dimensions of $[T]^{-1}$ and the corresponding star formation time-scales are given in the right-hand side ordinate. Symbols as in Fig. 10. Also plotted is the specific SFR required to double the stellar mass by $z = 0$ assuming a constant *specific* SFR, and the rate required to assemble the galaxy since the big bang assuming a constant SFR. Points lying above either line may be regarded as starbursting. Note that most of our detections can be regarded as starbursting. Right: corresponding predictions from the de Lucia et al. (2006) simulation. Note the obvious discrepancies with the observations.

Профили плотности темных гало: теория

- $\rho(r) = \rho_{cr} \delta_c / [(r/r_s)(1+r/r_s)^2]$

Navarro, Frenk, White (1996, 1997)

Профили плотности темной материи: наблюдения








Fig. 4.— Derived inner mass distributions from isotropic Jeans' equation analyses for six dSph galaxies. The modelling is reliable in each case out to radii of $\log(r)\text{kpc} \sim 0.5$. The unphysical behaviour at larger radii is explained in the text. The general similarity of the inner mass profiles is striking, as is their shallow profile, and their similar central mass densities. Also shown is an r^{-1} density profile, predicted by many CDM numerical simulations (eg Navarro, Frenk & White 1997).

Обзор KMOS-3D: $z=0.7-2.7$, панорамная спектроскопия на VLT



83% -Массивные, быстро вращающиеся диски, но ТОЛСТЫЕ И ГОРЯЧИЕ



Wisnioski et al. 2015

FALLING OUTER ROTATION CURVES OF STAR-FORMING GALAXIES AT $0.6 \lesssim Z \lesssim 2.6$ PROBED WITH KMOS^{3D} AND SINS/ZC-SINF

PHILIPP LANG^{1,2}, NATASCHA M. FÖRSTER SCHREIBER¹, REINHARD GENZEL^{1,3,4}, STIJN WUYTS⁵, EMILY WISNIOSKI¹, ALESSANDRA BEIFIORI^{1,6}, SIRIO BELL¹, RALF BENDER^{1,6}, GABE BRAMMER⁷, ANDREAS BURKERT^{1,6}, JEFFREY CHAN¹, RIC DAVIES¹, MATTEO FOSSATI^{1,6}, AUDREY GALAMETZ¹, SANDESH K. KULKARNI¹, DIETER LUTZ¹, J. TREVOR MENDEL¹, IVELINA G. MOMCHEVA⁷, THORSTEN NAAB⁸, ERICA J. NELSON¹, ROBERTO P. SAGLIA^{1,6}, STELLA SEITZ⁶, SANDRO TACCHELLA¹⁰, LINDA J. TACCONI¹, KEN-ICHI TADAKI¹, HANNAH ÜBLER¹, PIETER G. VAN DOKKUM⁹, DAVID J. WILMAN^{1,6}

¹Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse, D-85748 Garching, Germany

²Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

³Department of Physics, Le Conte Hall, University of California, Berkeley, CA 94720, USA

⁴Department of Astronomy, New Campbell Hall, University of California, Berkeley, CA 94720, USA

⁵Department of Physics, University of Bath, Claverton Down, Bath, BA2 7AY, UK

⁶Universitäts-Sternwarte München, Scheinerstr. 1, München, D-81679, Germany

⁷Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁸Max-Planck-Institut für Astrophysik, Karl Schwarzschildstr. 1, D-85748 Garching, Germany

⁹Astronomy Department, Yale University, New Haven, CT 06511, USA

¹⁰Institute of Astronomy, Department of Physics, Eidgenössische Technische Hochschule, ETH Zürich, CH-8093, Switzerland

101 галактика между $z=0.6$ и $z=2.6$ – просуммировали!



Table 1. Median properties of the stacking sample.

Property	Median
z	1.52
$\log(M_*[M_\odot])$	10.61
$\log(M_{\text{baryonic}} = M_* + M_{\text{gas}}^a)[M_\odot]$	10.87
$SFR [M_\odot/\text{yr}]$	41.9
$sSFR [\text{Gyr}^{-1}]$	1.20
$R_e^b [\text{kpc}]$	4.6
n^b	1.1

^aGas mass estimate based on empirical scaling relations.

^bIntrinsic effective radius along the major axis and Sérsic index derived from H -band Sérsic fits.

Показатель Серсика =1 → ЭКСПОНЕНЦИАЛЬНЫЕ ДИСКИ



... ВПИСАЛИ В КРИВУЮ ВРАЩЕНИЯ ИОНИЗОВАННОГО ГАЗА МОДЕЛЬ ЭКСПОНЕНЦИАЛЬНОГО ДИСКА

the noise cube associated with each data cube. Once we have constructed a rotation curve for each galaxy, we determine its observed amplitude and extent by fitting a Freeman exponential disk model of the form:

$$v_{\text{disk}}(r) = \frac{r}{r_d} \sqrt{\pi G \Sigma_0 r_d [I_0 K_0 - I_1 K_1]}, \quad (1)$$

where r_d is the radial scale-length of the exponential disk, corresponding to $r_d = R_e/1.68$. Σ_0 is the central mass surface density of the disk, and I_n and K_n denote the modified Bessel functions of the first and second kind (Freeman 1970). In the fitting, we leave r_d and Σ_0 as free parameters. Before fitting, the Freeman disk model is convolved with a 1D Gaussian of FWHM corresponding to the PSF associated with each galaxy. After the fit has converged, we determine the maximum observed velocity (V_{max}) and the radius of the peak (R_{turn}) from the model. In the right panels of Figure 2, rotation curves of two examples are shown together with the respective fit.




Figure 3. Final stacked pv diagram shown in logarithmic color scaling. The zero positions in velocity and radius are marked as black lines, and the radial position of $1R_{\text{turn}}$ and $2R_{\text{turn}}$ are marked as white dashed horizontal lines. The white symbols

Результаты






Figure 5. Top: Stacked rotation curve (black dots) plotted in units of normalized velocity (V/V_{max}), normalized radius (R/R_{turner}), and intrinsic effective radius (R/R_c). The error bars are derived from bootstrapping and include both sample variance as well as RMS noise in the spectra. The shaded area marks the half-light beam size of the average PSF observed for our sample. Middle: Effective number of galaxies contributing to the stack, accounting for masking out noisy pixels in the pv diagrams. The decrease in the number of contributing galaxies with increasing radius is driven by FOV limitations. Bottom: Median redshift and V_{rot}/σ_0 of contributing galaxies for a given radial bin.



Главный результат – темное гало НЕ чувствуется (но подразумевается...)



Genzel et al.: Nature!

Strongly baryon-dominated disk galaxies at the peak of galaxy formation ten billion years ago[†]

R.Genzel^{1,2*}, N.M. Förster Schreiber^{1*}, H.Übler¹, P.Lang¹, T.Naab³, R.Bender^{4,1}, L.J.Tacconi¹, E.Wisnioski¹, S.Wuyts^{1,5}, T.Alexander⁶, A. Beifiori^{4,1}, S.Belli¹, G. Brammer⁷, A.Burkert^{3,1}, C.M.Carollo⁸, J. Chan¹, R.Davies¹, M. Fossati^{1,4}, A.Galametz^{1,4}, S.Genel⁹, O.Gerhard¹, D.Lutz¹, J.T. Mendel^{1,4}, I.Momcheva¹⁰, E.J.Nelson^{1,10}, A.Renzini¹¹, R.Saglia^{1,4}, A.Sternberg¹², S.Tacchella⁸, K.Tadaki¹ & D.Wilman^{4,1}

¹Max-Planck-Institut für extraterrestrische Physik (MPE), Giessenbachstr.1, 85748 Garching, Germany
(genzel@mpe.mpg.de, forster@mpe.mpg.de)

²Departments of Physics and Astronomy, University of California, 94720 Berkeley, USA

³Max-Planck Institute for Astrophysics, Karl Schwarzschildstrasse 1, D-85748 Garching, Germany

⁴Universitäts-Sternwarte Ludwig-Maximilians-Universität (USM), Scheinerstr. 1, München, D-81679, Germany

⁵Department of Physics, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom

⁶Dept of Particle Physics & Astrophysics, Faculty of Physics, The Weizmann Institute of Science, POB 26, Rehovot 76100, Israel

⁷Space Telescope Science Institute, Baltimore, MD 21218, USA

⁸Institute of Astronomy, Department of Physics, Eidgenössische Technische Hochschule, ETH Zürich, CH-8093, Switzerland


⁹Center for Computational Astrophysics, 160 Fifth Avenue, New York, NY 10010, USA

¹⁰Department of Astronomy, Yale University, 260 Whitney Avenue, New Haven, CT 06511, USA

¹¹Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, Padova, I-35122, Italy

¹²School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

Выборка



Измерения




Figure 1. $H\alpha$ gas dynamics from KMOS and SINFONI in six massive star-forming galaxies. The galaxies have redshifts between $z=0.9$ and 2.4 . KMOS provides seeing-limited data (FWHM $\sim 0.6''$), and SINFONI allows both seeing-limited, and adaptive optics assisted observations (FWHM $\sim 0.2''$). For each galaxy column (a) shows the distribution of the integrated $H\alpha$ line surface brightness (with a linear colour scale), superposed on white contours of the stellar surface density (Σ_*) or the H-band continuum surface brightness (Σ_{H1}) (square root scaling). The vertical white bar denotes the

01351 and GS4 53501 we show both SINFONI (black filled circles) and KMOS (open blue circles) data sets, for D3a 15504 we show SINFONI data sets at $0.2''$ (black) and $0.5''$ (blue) resolution. Red continuous lines denote the best-fit dynamical model, constructed from a combination of a central compact bulge, an exponential disk and an NFW halo without adiabatic contraction, with a concentration of $c=4$ at $z\sim 2$ and $c=6.5$ at $z\sim 1$. For the modelling of the disk rotation, we also take into account the asymmetric drift correction inferred from the velocity dispersion curves ((d)-(e)¹⁴). Columns (d) and (e)

Еще измерения

Table 1. Physical Parameters of Observed Star-Forming Galaxies

	COS4 01351	D3a 6397	GS4 43501	zC 406690	zC 400569	D3a 15504
redshift	0.854	1.500	1.613	2.196	2.242	2.383
kpc/arcsec	7.68	8.46	8.47	8.26	8.23	8.14
Priors:						
M_* ($10^{11} M_\odot$)	0.54±0.16	1.2±0.37	0.41±0.12	0.42±0.12	1.2±0.37	1.1±0.34
$M_{\text{baryon}}(\text{gas+stars})$ ($10^{11} M_\odot$)	0.9±0.5	2.3±1.1	0.75±0.37	1.4±0.7	2.5±1.2	2.0±1.0
H-band $R_{1/2}$ (kpc)	8.6±1.3	5.9±0.8	4.9±0.7	5.5±1	4±2	6.3±1
inclination (°)	75±5	30±5	62±5	25±12	45±10	34±5
dark matter concentration parameter c	6.8	5	5	4	4	4
Fit parameters:						
$v_c(R_{1/2})$ (km/s) ^a	276	310	257	301	364	299
$R_{1/2}(n=1)$ (kpc)	7.3	7.4	4.9	5.5	3.3	6
σ_0 (km/s)	39	73	39	74	34	76
$M_{\text{baryon}}(\text{gas+stars, including bulge})$ ($10^{11} M_\odot$)	1.7	2.3	1.0	1.7	1.7	2.1
$M_{\text{bulge}}/M_{\text{baryon}}$	0.2	0.35	0.4	0.6	0.37	0.15
$f_{\text{DM}}(R_{1/2})=(v_{\text{DM}}/v_c)^2 _{R=R_{1/2}}$ ^b	0.21 (±0.1)	0.17 (<0.38)	0.19 (±0.09)	0.0 (<0.08)	0.0 (<0.07)	0.12 (<0.26)

^a Total circular velocity at the half-light radius (rest-frame optical) $R_{1/2}$, including bulge, exponential disk ($n=1$) and dark matter, and corrected for asymmetric drift: $v_c(R)^2 = v_{\text{rot}}(R)^2 + 3.36 \sigma_0^2 \times (R/R_{1/2})|_{n=1}$.

^b Ratio of dark matter to total mass at the half-light radius of the optical light, $f_{\text{DM}}(R_{1/2}) = (v_{\text{DM}}/v_c)^2|_{R=R_{1/2}}$, with numbers in the parentheses giving the ± 2 rms ($\delta\chi^2=4$, ~95% probability) uncertainties, or upper limits. We use an NFW halo of concentration parameter c, and no adiabatic contraction.

Главный результат: отсутствие темной материи – признак раннего типа галактики?




Figure 2. Normalized rotation curves. (a): The various symbols denote the folded and binned rotation curve data for the six galaxies in Figure 1, combined with the stacked rotation curve of 97 $z=0.6-2.6$ star-forming galaxies¹⁸ (Methods). For all rotation curves we averaged data points located symmetrically on either side of the dynamical centres, and plot the rotation velocities and radii normalized to their maximum values. Error bars are ± 1 rms. (b): The black data points




Figure 3. Dark matter fractions. Dark matter fractions from different methods are listed as a function of the circular velocity of the disk, at the half mass/light radius of the disk, for galaxies in the current Universe and ~ 10 Gyr ago. The large blue circles with red outlines indicate the dark matter fractions derived from the outer-disk rotation curves of the six high- z disks presented in this paper (Table 1), along with the ± 2 rms

Выводы и спекуляции:

- На $z=2$ (10 млрд лет назад) в дисковых галактиках в пределах оптического радиуса доминируют барионы – в среднем до 90% общей динамической массы.
- С уменьшением z – с ходом эволюции Вселенной – доля барионов внутри оптического радиуса дисковых галактик ПЛАВНО падает.
- А на $z=0$ в пределах оптического радиуса барионы доминируют только в массивных галактиках ранних типов – РОДСТВЕННАЯ СВЯЗЬ? ЭФФЕКТ МАССЫ? ДРУГАЯ ПРИРОДА СОВРЕМЕННОГО ЗВЕЗДООБРАЗОВАНИЯ?